

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Diisopropylammonium hydrogen phenylphosphonate

Modou Sarr,^a Mouhamadou Sembene Boye,^{a*} Aminata Diasse-Sarr,^a Arnaud Grosjean^b and Philippe Guionneau^b

^aLaboratoire de Chimie Minérale et Analytique, Département de Chimie, Faculté des Sciences et Techniques, Université Cheikh Anta Diop, Dakar, Senegal, and ^bCNRS, Université de Bordeaux, ICMCB, 87 Avenue du Dr A. Schweitzer, Pessac F-33608, France

Correspondence e-mail: mouhasboye@hotmail.com


Received 17 September 2012; accepted 30 September 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.029; wR factor = 0.080; data-to-parameter ratio = 16.4.

In the title salt, $[(\text{CH}_3)_2\text{CH}]_2\text{NH}_2^+ \cdot [\text{C}_6\text{H}_5\text{PO}_2(\text{OH})]^-$, the anions are linked by pairs of $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds, forming inversion dimers. These dimers are bridged by the cations *via* $\text{N}-\text{H} \cdots \text{O}$ hydrogen bonds, leading to a three-dimensional structure.

Related literature

For crystal structures of closely related compounds, see: Diop *et al.* (2012); Beckmann *et al.* (2003).



Experimental

Crystal data

$\text{C}_6\text{H}_{16}\text{N}^+ \cdot \text{C}_6\text{H}_5\text{O}_3\text{P}^-$

$M_r = 259.28$

Monoclinic, $P2_1/n$
 $a = 11.9166$ (2) Å
 $b = 9.0982$ (1) Å
 $c = 12.8539$ (1) Å
 $\beta = 101.013$ (1)°
 $V = 1367.95$ (3) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.20$ mm⁻¹
 $T = 293$ K
 $0.88 \times 0.63 \times 0.25$ mm

Data collection

Nonius KappaCCD diffractometer
 Absorption correction: multi-scan
 (*SCALEPACK*; Otwinowski & Minor, 1997)
 $T_{\min} = 0.845$, $T_{\max} = 0.952$

7738 measured reflections
 3990 independent reflections
 3626 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.013$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$
 $wR(F^2) = 0.080$
 $S = 1.03$
 3990 reflections

243 parameters
 All H atoms refined
 $\Delta\rho_{\max} = 0.38$ e Å⁻³
 $\Delta\rho_{\min} = -0.33$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{N1}-\text{H17} \cdots \text{O3}$	0.900 (14)	1.960 (14)	2.8510 (10)	170.3 (12)
$\text{O2}-\text{H6} \cdots \text{O3}^{\text{i}}$	0.847 (19)	1.744 (19)	2.5895 (10)	177.4 (19)
$\text{N1}-\text{H16} \cdots \text{O1}^{\text{ii}}$	0.916 (15)	1.764 (15)	2.6782 (10)	176.7 (13)

Symmetry codes: (i) $-x + 2, -y + 1, -z + 2$; (ii) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *COLLECT* (Hooft, 1998); cell refinement: *HKL DENZO* (Otwinowski & Minor, 1997); data reduction: *SCALEPACK* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *publCIF* (Westrip, 2010).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2589).

References

- Beckmann, J., Dakternieks, D. & Duthie, A. (2003). *Appl. Organomet. Chem.* **17**, 817–818.
 Diop, T., Diop, L., Maris, T. & Stoeckli-Evans, H. (2012). *Acta Cryst.* **E68**, o1432.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Hooft, R. (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
 Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2012). E68, o3078 [doi:10.1107/S1600536812041086]

Diisopropylammonium hydrogen phenylphosphonate

Modou Sarr, Mouhamadou Sembene Boye, Aminata Diasse-Sarr, Arnaud Grosjean and Philippe Guionneau

Comment

In the asymmetric unit of the title salt (Fig. 1), the anion, hydrogen phenylphosphonate (PhPO_3H^-), adopts a tetrahedral geometry, with three oxygen atoms and a benzene group with O—P—O and O—P—C angles in the ranges $109.58(4) - 115.20(4)$ and $105.24(4) - 108.51(4)^\circ$, respectively. Two P—O distances in the anion are close, ($\text{P1—O1} = 1.4932(7)$ and $\text{P1—O3} = 1.5191(7)$ Å) indicating the presence of extensive π -delocalization of the P=O double bonds. The bond distance P1—O2 ($1.5809(7)$ Å for P—OH bond) is significantly longer than the other two P—O bonds. The P—O bond distances in the title salt agree very well with the corresponding bond distances reported in closely related compounds (Diop *et al.*, 2012; Beckmann *et al.*, 2003).

In the crystal, the anions are connected by $\text{O2—H6}\cdots\text{O3}$ hydrogen bonds forming dimers about inversion centers. These pairs are then bridged through cations *via* $\text{N—H}\cdots\text{O}$ hydrogen bonds leading to a three-dimensional structure (Tab. 1 & Fig. 2).

Experimental

The title compound was synthesized by mixing $[(\text{CH}_3)_2\text{CH}]_2\text{NH}$ and PhPO_3H_2 in water (1/1 ratio). The precipitate obtained was filtered. The crystals suitable for X-ray crystallographic analysis were grown from a solution of water by slow evaporation at room temperature.

Refinement

All H atoms were located from difference maps and were allowed to refine freely with U_{iso} .

Computing details

Data collection: *COLLECT* (Hooft, 1998); cell refinement: *HKL DENZO* (Otwinowski & Minor, 1997); data reduction: *SCALEPACK* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *pubCIF* (Westrip, 2010).




Figure 1

The asymmetric unit of the title salt. Displacement ellipsoids are drawn at the 50% probability level.




Figure 2

A view of the O—H···O and N—H···O hydrogen bonds (light-blue colored lines) in the crystal structure of the title compound. H atoms non-participating in hydrogen-bonding were omitted for clarity.

Diisopropylammonium hydrogen phenylphosphonate

Crystal data

$\text{C}_6\text{H}_{16}\text{N}^+\cdot\text{C}_6\text{H}_5\text{O}_3\text{P}^-$

$M_r = 259.28$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1n$

$a = 11.9166\ (2)\ \text{\AA}$

$b = 9.0982\ (1)\ \text{\AA}$

$c = 12.8539\ (1)\ \text{\AA}$

$\beta = 101.013\ (1)^\circ$

$V = 1367.95\ (3)\ \text{\AA}^3$

$Z = 4$

$F(000) = 560$

$D_x = 1.259\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 4223 reflections

$\theta = 0.4\text{--}30.0^\circ$

$\mu = 0.20\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Prism, colourless

$0.88 \times 0.63 \times 0.25\ \text{mm}$

Data collection

Nonius KappaCCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ scans, and ω scans with κ

Absorption correction: multi-scan

(*SCALEPACK*; Otwinowski & Minor, 1997)

$T_{\min} = 0.845$, $T_{\max} = 0.952$

7738 measured reflections
3990 independent reflections
3626 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.013$

$\theta_{\text{max}} = 30.1^\circ$, $\theta_{\text{min}} = 3.2^\circ$
 $h = -16 \rightarrow 16$
 $k = -12 \rightarrow 12$
 $l = -18 \rightarrow 18$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.029$
 $wR(F^2) = 0.080$
 $S = 1.03$
3990 reflections
243 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods
Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites
All H-atom parameters refined
 $w = 1/[\sigma^2(F_o^2) + (0.0397P)^2 + 0.4747P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.38 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.33 \text{ e } \text{\AA}^{-3}$
Extinction correction: *SHELXL*,
 $\text{Fc}^* = k\text{Fc}[1 + 0.001 \times \text{Fc}^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Extinction coefficient: 0.013 (2)

Special details

Experimental. The analytical data - % calculated (% found): C: 55.59 (55.64); H: 8.55 (8.91); N: 5.40 (5.58).
Infrared data (cm^{-1}) [br= broad; s= strong; m=medium]
2694br νNH ; 1148m, 1129 s, 1059, 1027m νPO_3 ; 898 s δPO_3 ; 752 m νPC ; 708 s, 696 s νPh .
Melting point = 437–438 K

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
P1	0.966159 (19)	0.40247 (2)	0.846319 (17)	0.01271 (8)
O2	1.04330 (6)	0.32302 (7)	0.94386 (5)	0.01729 (14)
N1	0.66048 (7)	0.51553 (9)	0.78554 (6)	0.01430 (15)
C1	1.06532 (7)	0.49048 (10)	0.77593 (7)	0.01453 (17)
C6	1.14485 (9)	0.59178 (11)	0.82778 (8)	0.0222 (2)
C12	0.62928 (8)	0.41946 (10)	0.68876 (8)	0.01742 (18)
C13	0.69808 (9)	0.47210 (12)	0.60787 (8)	0.0225 (2)
C2	1.06337 (9)	0.46114 (11)	0.66912 (8)	0.01979 (19)
C4	1.21864 (9)	0.63037 (13)	0.66783 (9)	0.0270 (2)
C3	1.13968 (9)	0.53095 (13)	0.61530 (9)	0.0256 (2)
C5	1.22118 (10)	0.66111 (13)	0.77426 (10)	0.0287 (2)
O1	0.89926 (6)	0.29081 (8)	0.77497 (5)	0.02025 (15)
C8	0.60257 (8)	0.48352 (11)	0.87717 (8)	0.01757 (18)
C9	0.64492 (9)	0.59615 (12)	0.96314 (8)	0.0215 (2)
C11	0.50164 (9)	0.42810 (13)	0.64537 (9)	0.0245 (2)
C10	0.62574 (10)	0.32619 (12)	0.91508 (9)	0.0257 (2)

O3	0.89542 (5)	0.52144 (7)	0.88605 (5)	0.01657 (14)
H6	1.0624 (16)	0.376 (2)	0.9984 (15)	0.054 (5)*
H17	0.7362 (12)	0.5109 (14)	0.8110 (11)	0.022 (3)*
H16	0.6429 (12)	0.6104 (16)	0.7649 (11)	0.028 (3)*
H21	0.5218 (11)	0.4980 (14)	0.8513 (10)	0.018 (3)*
H18	0.6510 (11)	0.3197 (15)	0.7094 (10)	0.020 (3)*
H10	0.6767 (12)	0.5719 (16)	0.5864 (11)	0.029 (4)*
H7	0.7279 (12)	0.5857 (15)	0.9907 (11)	0.027 (3)*
H12	0.6818 (13)	0.4116 (16)	0.5460 (12)	0.034 (4)*
H9	0.6088 (13)	0.5821 (17)	1.0245 (12)	0.036 (4)*
H11	0.7788 (13)	0.4705 (16)	0.6362 (12)	0.032 (4)*
H8	0.6298 (13)	0.6961 (18)	0.9372 (12)	0.035 (4)*
H13	0.4820 (12)	0.3727 (17)	0.5783 (12)	0.034 (4)*
H20	0.7076 (13)	0.3059 (16)	0.9306 (11)	0.029 (3)*
H19	0.5963 (13)	0.3128 (18)	0.9806 (13)	0.041 (4)*
H15	0.4790 (13)	0.5297 (17)	0.6277 (12)	0.034 (4)*
H14	0.4571 (13)	0.3869 (16)	0.6953 (12)	0.034 (4)*
H22	0.5884 (13)	0.2571 (18)	0.8637 (13)	0.038 (4)*
H5	1.1462 (12)	0.6131 (15)	0.9024 (11)	0.028 (3)*
H3	1.2709 (13)	0.6789 (17)	0.6323 (12)	0.037 (4)*
H4	1.2746 (13)	0.7291 (18)	0.8131 (12)	0.040 (4)*
H2	1.1388 (13)	0.5080 (17)	0.5413 (12)	0.034 (4)*
H1	1.0101 (12)	0.3924 (16)	0.6328 (11)	0.027 (3)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
P1	0.01307 (12)	0.01214 (12)	0.01264 (12)	−0.00065 (7)	0.00175 (8)	−0.00020 (7)
O2	0.0219 (3)	0.0150 (3)	0.0141 (3)	0.0025 (2)	0.0012 (2)	0.0012 (2)
N1	0.0134 (3)	0.0143 (3)	0.0154 (3)	0.0003 (3)	0.0031 (3)	0.0005 (3)
C1	0.0141 (4)	0.0135 (4)	0.0161 (4)	0.0019 (3)	0.0032 (3)	0.0015 (3)
C6	0.0229 (5)	0.0226 (5)	0.0214 (5)	−0.0063 (4)	0.0051 (4)	−0.0012 (4)
C12	0.0184 (4)	0.0157 (4)	0.0179 (4)	0.0004 (3)	0.0027 (3)	−0.0029 (3)
C13	0.0214 (5)	0.0297 (5)	0.0171 (4)	0.0020 (4)	0.0049 (4)	−0.0015 (4)
C2	0.0207 (4)	0.0222 (5)	0.0171 (4)	0.0005 (4)	0.0052 (3)	−0.0001 (3)
C4	0.0241 (5)	0.0279 (5)	0.0321 (6)	0.0004 (4)	0.0129 (4)	0.0102 (4)
C3	0.0275 (5)	0.0309 (6)	0.0209 (5)	0.0033 (4)	0.0110 (4)	0.0049 (4)
C5	0.0262 (5)	0.0280 (5)	0.0326 (6)	−0.0105 (4)	0.0077 (4)	0.0014 (4)
O1	0.0231 (3)	0.0169 (3)	0.0193 (3)	−0.0056 (3)	0.0003 (3)	−0.0021 (3)
C8	0.0148 (4)	0.0209 (4)	0.0182 (4)	0.0006 (3)	0.0064 (3)	0.0026 (3)
C9	0.0228 (5)	0.0247 (5)	0.0179 (4)	0.0034 (4)	0.0065 (4)	−0.0007 (4)
C11	0.0190 (4)	0.0267 (5)	0.0263 (5)	−0.0035 (4)	0.0005 (4)	−0.0067 (4)
C10	0.0306 (5)	0.0215 (5)	0.0264 (5)	−0.0017 (4)	0.0089 (4)	0.0066 (4)
O3	0.0137 (3)	0.0197 (3)	0.0161 (3)	0.0026 (2)	0.0023 (2)	−0.0017 (2)

Geometric parameters (\AA , $^\circ$)

P1—O1	1.4932 (7)	C2—C3	1.3962 (14)
P1—O3	1.5191 (7)	C2—H1	0.948 (14)
P1—O2	1.5809 (7)	C4—C3	1.3845 (17)

P1—C1	1.8068 (9)	C4—C5	1.3910 (16)
O2—H6	0.847 (19)	C4—H3	0.949 (15)
N1—C8	1.5030 (12)	C3—H2	0.972 (15)
N1—C12	1.5077 (12)	C5—H4	0.957 (16)
N1—H17	0.900 (14)	C8—C10	1.5202 (14)
N1—H16	0.916 (15)	C8—C9	1.5208 (14)
C1—C2	1.3945 (13)	C8—H21	0.965 (13)
C1—C6	1.3966 (13)	C9—H7	0.989 (15)
C6—C5	1.3920 (14)	C9—H9	0.976 (15)
C6—H5	0.976 (14)	C9—H8	0.973 (16)
C12—C11	1.5188 (14)	C11—H13	0.987 (15)
C12—C13	1.5193 (14)	C11—H15	0.977 (16)
C12—H18	0.967 (13)	C11—H14	0.982 (15)
C13—H10	0.969 (15)	C10—H20	0.976 (15)
C13—H12	0.957 (15)	C10—H19	0.980 (16)
C13—H11	0.961 (15)	C10—H22	0.957 (16)
O1—P1—O3	115.20 (4)	C3—C4—C5	119.74 (9)
O1—P1—O2	109.72 (4)	C3—C4—H3	121.1 (9)
O3—P1—O2	109.58 (4)	C5—C4—H3	119.1 (9)
O1—P1—C1	108.51 (4)	C4—C3—C2	120.12 (10)
O3—P1—C1	108.10 (4)	C4—C3—H2	119.8 (9)
O2—P1—C1	105.24 (4)	C2—C3—H2	120.1 (9)
P1—O2—H6	114.8 (12)	C4—C5—C6	120.10 (10)
C8—N1—C12	117.16 (7)	C4—C5—H4	122.1 (10)
C8—N1—H17	106.5 (8)	C6—C5—H4	117.8 (10)
C12—N1—H17	110.1 (8)	N1—C8—C10	110.56 (8)
C8—N1—H16	107.3 (9)	N1—C8—C9	107.49 (8)
C12—N1—H16	107.5 (9)	C10—C8—C9	112.76 (9)
H17—N1—H16	107.9 (12)	N1—C8—H21	106.3 (7)
C2—C1—C6	118.58 (9)	C10—C8—H21	110.3 (8)
C2—C1—P1	121.24 (7)	C9—C8—H21	109.2 (7)
C6—C1—P1	120.17 (7)	C8—C9—H7	111.5 (8)
C5—C6—C1	120.74 (10)	C8—C9—H9	111.2 (9)
C5—C6—H5	120.4 (8)	H7—C9—H9	105.2 (12)
C1—C6—H5	118.8 (8)	C8—C9—H8	111.5 (9)
N1—C12—C11	110.17 (8)	H7—C9—H8	108.6 (12)
N1—C12—C13	107.55 (8)	H9—C9—H8	108.5 (12)
C11—C12—C13	111.46 (9)	C12—C11—H13	110.3 (8)
N1—C12—H18	107.9 (8)	C12—C11—H15	110.3 (9)
C11—C12—H18	110.2 (8)	H13—C11—H15	105.8 (12)
C13—C12—H18	109.5 (7)	C12—C11—H14	111.6 (9)
C12—C13—H10	110.1 (8)	H13—C11—H14	108.0 (12)
C12—C13—H12	109.8 (9)	H15—C11—H14	110.7 (12)
H10—C13—H12	107.5 (12)	C8—C10—H20	110.9 (8)
C12—C13—H11	111.6 (9)	C8—C10—H19	108.6 (9)
H10—C13—H11	108.2 (12)	H20—C10—H19	108.0 (12)
H12—C13—H11	109.6 (12)	C8—C10—H22	111.4 (9)
C1—C2—C3	120.72 (10)	H20—C10—H22	109.3 (12)

C1—C2—H1	119.7 (8)	H19—C10—H22	108.3 (13)
C3—C2—H1	119.6 (8)		
O1—P1—C1—C2	6.07 (9)	C8—N1—C12—C13	−179.89 (8)
O3—P1—C1—C2	−119.51 (8)	C6—C1—C2—C3	0.34 (14)
O2—P1—C1—C2	123.46 (8)	P1—C1—C2—C3	179.35 (8)
O1—P1—C1—C6	−174.93 (8)	C5—C4—C3—C2	−0.39 (17)
O3—P1—C1—C6	59.48 (8)	C1—C2—C3—C4	0.11 (16)
O2—P1—C1—C6	−57.55 (8)	C3—C4—C5—C6	0.21 (18)
C2—C1—C6—C5	−0.53 (15)	C1—C6—C5—C4	0.26 (17)
P1—C1—C6—C5	−179.55 (8)	C12—N1—C8—C10	58.07 (11)
C8—N1—C12—C11	58.44 (11)	C12—N1—C8—C9	−178.48 (8)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H17 \cdots O3	0.900 (14)	1.960 (14)	2.8510 (10)	170.3 (12)
O2—H6 \cdots O3 ⁱ	0.847 (19)	1.744 (19)	2.5895 (10)	177.4 (19)
N1—H16 \cdots O1 ⁱⁱ	0.916 (15)	1.764 (15)	2.6782 (10)	176.7 (13)

Symmetry codes: (i) $-x+2, -y+1, -z+2$; (ii) $-x+3/2, y+1/2, -z+3/2$.